

RINSHO HINYOKIKA

(JAPANESE JOURNAL OF CLINICAL UROLOGY)

Advisory Board : **Toyohei Machida MD, Akimi Ogawa MD, Masao Akimoto MD, Masaru Murai MD, Kenjiro Kohri MD, Tomoaki Fujioka MD**

Editorial Board : **Mototsugu Oya MD, Yukihiro Kondo MD, Yoshiyuki Kojima MD**

CONTENTS

Vol 68 No 9 August 2014

SPECIAL THEME : Sudden change of a patient in urology—How do you evaluate and cope?

- When a lot of urine drainage was performed, the patient lost consciousness —Tetsuya Takao 640
- After injecting LH-RH analogue, subcutaneous hemorrhage
was seen in the abdominal injection site —Takanobu Utsumi, et al 644
- Emergency in urological test :
- Blood pressure drops rapidly when local or caudal anesthesia is done —Yuki Kitamura, et al 648
- The shock measures on the cystoscopy —So Omori 652
- The blood pressure of the patient fell after the administration of contrast
medium for urography and he has begun to complain of dyspnea —Koichi Nakajima 654
- Management for the oxygen desaturation occurring soon after the start
of transurethral resection or transurethral ureterolithotripsy —Hironori Betsunoh 658
- A patient presenting with yawn followed by nausea, vomiting,
and lastly agitated confusional states during transurethral resection
or ureteroscopy —Yasuo Kohjimoto 662
- Management of sudden circulatory collapse during transurethral surgery —Shuzo Hamamoto 666
- After both open and laparoscopic surgeries, patients' blood pressures
have dropped suddenly and pulse rates have increased —Hidetoshi Akita 670
- A patient complained of severe abdominal pain after an open
or an endoscopic surgery —Norihiro Tsuchiya 674
- Drain drainage of patients has increased suddenly after open surgery
and laparoscopic surgery —Yukihiro Umemoto 678
- Evaluation of postoperative oliguria —Yuji Tokuda 680
- A patient experienced acute dyspnea and chest pain at first
postoperative walking after a open or laparoscopic surgery —Shingo Hatakeyama, et al 685
- Emergency management for the patient with low blood pressure
after chemotherapy or molecular targeted therapy —Fumiya Hongo, et al 688
- Dyspnea during chemotherapy or molecular targeting therapy —Noboru Nakaigawa 692
- Fever in cancer patients treated with anti-cancer agents —Takashi Karashima, et al 696
- A sudden change after the treatment with anti-cancer drugs :
- Neurological symptoms such as the disturbance of consciousness
and convulsions —Mitsuru Noguchi, et al 700