

Original Article

- Development of a transparent drape to observe intraoperative positioning in laparoscopic surgery
—Comprehensive assessment in the medical professions and patients —Eriko KAWASAKI, et al ● 245

Case Report

- A case of pelvic accessory spleens which were difficult to diagnose preoperativelyJun KONO, et al ● 251
- Elective surgery of totally extraperitoneal approach after manual reduction of incarcerated obturator herniaReina MIYAMOTO, et al ● 257
- Pure laparoscopic partial caudate hepatectomy for hepatocellular carcinomaKatsuhiko ITO, et al ● 263
- A case of laparoscopic removal of translocated intrauterine device from peritoneal cavityNatsuki KOIDE, et al ● 269
- A case of unruptured aneurysm of cystic artery treated by laparoscopic cholecystectomyHisashi NISHIDA, et al ● 275
- A case of laparoscopic surgery for ileus caused by misplacement of percutaneous endoscopic gastrostomy passed through the mesentery of the small bowelSoichi TAGAMI, et al ● 281
- Laparoscopic cholecystectomy in an adult male with Duchenne muscular dystrophyYusuke YAMAOKA, et al ● 287
- A case of laparoscopic retroperitoneal drainage for retroperitoneal abscess due to idiopathic duodenal perforation in a patient on maintenance hemodialysisTadashi SAKURAI, et al ● 293
- A case of cholecystogastric fistula treated by laparoscopic surgeryHiroki SATO, et al ● 297
- Transverse mesocolon hernia treated by laparoscopic surgery : a case reportTadahiro KAMIYA, et al ● 305
- A minute tumor of the small intestine resected by intraoperative CT-angiography and ICG fluorescent imaging techniques : a case reportToshifumi WATANABE, et al ● 311

Operative Technique

- Robot-assisted thoracoscopic esophagectomy with the patient in the prone positionYoshiaki OSAKA, et al ● 317
- A novel and safe technique for esophagojejunostomy using the direct inserted anvil after laparoscopy assisted total gastrectomyRyusuke AIHARA, et al ● 323
- Reduced port surgery (RPS) for left-sided colorectal disease with transanal extraction of the specimen performed with a maximum incision wound of 5 mmEiichiro TOYAMA, et al ● 329