

Diagnostic Imaging in the Hepatobiliary and Pancreatic Regions : Focusing on the Differentiation between Benign and Malignant Lesions

■ Foreword	Takamichi Murakami et al	637
■ Differentiation between benign and malignant hepatocellular nodule during multistep hepatocarcinogenesis	Satoshi Kobayashi et al	639
■ MR imaging diagnosis of hepatocellular adenoma : imaging features, differential diagnosis, and recent topics	Hiroki Haradome	649
■ CT/MR diagnosis of biliary system : differentiation between benign and malignant lesions	Kengo Yoshimitsu et al	663
■ Tips and pitfalls in the differentiation between benign and malignant solid lesions of the pancreas	Yasunari Fujinaga et al	681
■ Differentiation between benign and malignancy in pancreatic cystic lesions by using imaging modalities	Shunro Matsumoto et al	691
* * *		
■ A practical technique for the postoperative radiotherapy in breast cancer (3) : technique for large field matching	Kenji Sekiguchi et al	703
■ HRCT features of interstitial lung disease patients with antisynthetase syndrome	Yoshida Atsushi et al	715
■ Correlation of ¹⁸ F FDG-PET/CT findings with abdominal aortic aneurysm growth	Mami Morimoto et al	721
■ Effective occlusion with Embosphere® before extraction in gingival hemangioma	Yusuke Arai et al	727
■ A case of cervical actinomycosis	Ryoya Ochiai et al	733
■ A case of pleomorphic adenoma of the breast	Takeshi Oshima et al	739
■ A case of ovarian cystadenofibroma with atypical presentation on MR imaging	Akira Baba et al	745
■ Efficacy of RT for Intradural Hemangiopericytomas repeated recurrence : a case report	Tamaki Kondo et al	749

Editors

Yutaka Imai (Tokai University School of Medicine)
Takehiko Gokan (Showa University School of Medicine)
Yasuyuki Kurihara (St. Luke's International Hospital)
Jun Hashimoto (Tokai University School of Medicine)
Harushi Mori (The University of Tokyo Graduate School of Medicine)
Katsumasa Nakamura (Hamamatsu University School of Medicine)